

Toxic effects of Tebufenozide on human sperm function *in vitro*

¹Pavithra D. Dayananda

¹Dinithi C Peiris (Corresponding author)

¹Department of Zoology, University of Sri
Jayewardenepura

Gagodawila, Nugegoda 1020, Sri Lanka

Abstract

Tebufenozide is a widely used insecticide in agriculture specially against lepidopteron, but their effects in the male reproductive tract are scarcely known. Hence, the present study has evaluated the effects of tebufenozide on human sperm function *in vitro*. Human sperm were incubated with equivalent doses of 0 (control; BWW medium), 1.25 (low dose), 2.5 (mid dose), 5 (high dose) $\mu\text{g/ml}$ and total motility and sperm viability was evaluated. Similarly, sperm capacitation, acrosome reaction and sperm DNA damage was examined. Result showed significant declining of total sperm motility at all doses (1.25 $\mu\text{g/ml}$: by 33.7%; 2.5 $\mu\text{g/ml}$ by 36.4%; 5 $\mu\text{g/ml}$: 36.3%) and sperm viability was reduced significantly only at the highest dose by 46.2%. DNA damages were significantly increased by 164.8% with 1.25 $\mu\text{g/ml}$ dose, by 326.9% with 2.5 $\mu\text{g/ml}$ dose and by 357.9% with 5 $\mu\text{g/ml}$ dose. Percentage of capacitated spermatozoa were affected at all dose levels (by 20.6%: low dose; 45.5%: mid dose and 58.0%: high dose), but acrosome reaction was impaired only at the highest dose (36.6%). The observations suggest that exposure to tebufenozide and its accumulation in the seminal plasma may increase the risk of infertility.

KEYWORDS: Tebufenozide, growth hormone regulator, reproductive toxicant, sperm function

1. Introduction

Use of environmental chemicals have increased as a consequence of efforts to meet increase food demand. Pesticides are potentially hazards to individuals involved in manufacture, formulation and applications in the field (Sankoh et al., 2016). Due to increase use of pesticides, there has been a substantial increase in the number of pesticides in the environment (Sharma et al. 2005). Studies have shown that increase usage of man-made chemicals result in destructive reproductive function in animals (Peiris et al. 1995 and Peiris-John & Wickremasinghe, 2008) and in humans (Skakkebaek et al., 2006). It is known that chemical compounds binds to receptors and acts either as an agonist or antagonist (Luconi et al., 2001) and could induce DNA damages in human spermatozoa (Kudavidanage and Peiris, 2016).

Tebufenozoid (trade name mimic 20F) is a widely used insecticides in Sri Lanka. It is an insecticide which, acts as a hormone that induces metamorphosis of insect larvae (Dharmadasa et al., 2008). An association between exposure to insecticides and reproductive disorders such as infertility, birth defect, adverse pregnancy outcomes and perinatal deaths has been shown by several authors (Peiris and Moore 2001a, Peiris and Moore 2001b, Ratnasooriya et al. 1996). Pesticides are known to influence the gonads and to cause abnormal semen parameters with decreased sperm concentration (Perry et al., 2007), decreased sperm volume and decreased sperm counts (Recio-Vega *et al.*, 2008). Once spermatozoa are produced in the testes, they undergo maturation in the epididymis. Upon ejaculation, in the female reproductive tract, spermatozoa undergo a process known as capacitation. To penetrate the egg zona pelucida, the sperm releases its enzyme via a process known as acrosome reaction. Any disturbances in any of these processes will impair sperm function.

Although effects of these pesticides on other animals were recorded the effects on human is limited. Hence the present aimed to determine the effects of tebufenozoid on sperm motility, sperm vitality sperm capacity and acrosome reaction *in vitro*.

2. Materials and methods

2.1 Test material

Unformulated tebufenozoid (mimic 20F, purity: 99.0%) was obtained from Haylee's Agriculture Ltd., Colombo, Sri Lanka.

2.2 Semen collection and preparation

Semen samples were collected from healthy male (age 20 - 28 years) donors from University of Sri Jayewardenepura, Sri Lanka in a sterile specimen vial. Prior to semen collection donors were given an information sheet and a consent form seeking their willingness to participate in the study. All participating subjects were asked to abstain from any sexual activity for 3 to 5 days before semen collection. After liquefaction at 37°C for 30 min, semen quality parameters were measured (n=9) according to World Health Organization guidelines (WHO, 2010). Only ejaculates from healthy volunteers with normal parameters (sperm concentration > 40 x 10⁶ spermatozoa/ml, total sperm motility > 50%, normal sperm morphology > 50%) were included in this study. Ethical approval was obtained by Ethics Review Committee, Faculty of Medical Sciences, University of Sri Jayewardenepura, Nugegoda, Sri Lanka

2.3 Experimental design

The doses were selected according to the field recommended dose for tebufenozide. The LD₅₀ values of field recommended doses were selected the highest dose levels while one half of and one fourth of the LD₅₀ values were taken as mid and the lowest dose levels respectively. However, to compare the toxicity of tebufenozide, the same dosage (low dose: 1.25 µg/ml, mid dose: 2.5 µg/ml and high dose: 5 µg/ml) were used for the study. All the insecticides are readily dissolved in water and hence, Biggers Whitten Whittingham (BWW) media was taken as the control. Sperm suspensions (fixed to 40 × 10⁶ spermatozoa/ml) were placed in Eppendorf tubes and desired volume of either isotonic saline or test solution were added (final volume of the tube is equal to 1 ml).

2.4 Assessment of total sperm motility

Samples were mixed with different concentrations (1.25 µg/ml, 2.5 µg/ml, 5 µg/ml) of tebufenozide or BWW (control) and sperm suspensions were then incubated in a humidified incubator (Sanyo Electric Co. Ltd, Japan) at 37°C in 5% CO₂ for 4h. Subsequently, total percentage motile spermatozoa (WHO, 2010) were estimated (by counting approximately 100 cells) under Olympus microscope with phase contrast optics (X 400; Olympus Corporation, Japan).

2.5 Assessment of sperm viability

The viability of spermatozoa of treated with different doses of tebufenozide (n=6) and the control were assessed using Eosin Y stain technique (WHO, 2010) upon incubation for 4 h. Spermatozoa with red or pink colouration in their head regions was classified as 'dead' and those appearing white or light pink were classified as 'live' and percentage vitality was calculated. $\text{Percentage vitality} = (\text{total number of cells} - \text{number stained} / \text{total number of cells}) \times 100$. Total number of cells = 100.

2.6 Assessment of sperm capacitation

Two hundred μl of BWW medium was placed at the center of a glass Petri dish and covered with liquid paraffin oil (BDH Chemical Ltd., Poole, UK). Subsequently, 100 μl of sperm suspension was added to the BWW medium in the petri dish and mixed well. Either 100 μl of different doses of tebufenozide (50, 100, 200 $\mu\text{g/ml}$) or BWW (control) were added to the BWW medium in a Petri dish (n=6). These were incubated at 37 °C for 3h in 5% CO₂ and 95% O₂. Subsequently 10 μl of each of these sperm suspensions was transferred on to a warm glass slide at 37 °C and was examined using phase contrast microscope (Olympus Corporation, Japan) at 100 magnifications. Then the number of hyper activated sperms and totally immotile sperms were recorded (at least 100 spermatozoa per suspension). A spermatozoon was considered to be hyper activated if it followed a nonlinear swimming with vigorous tail movement and marked lateral excursions of head.

2.7 Assessment of acrosome reaction

Sperm ability to undergo acrosome reaction was determined according to the method described by Gupta et al. (1993). Briefly, the acrosome reaction was induced by incubating spermatozoa with calcium ionophore A23187 with three doses of tebufenozide and the control (n=6). After 3 h incubation at 37 °C for 3h in 5% CO₂ incubator, sperm samples were centrifuged and acrosomal status were determined using fluoresceinated *Pisum sativum* agglutinin. Sperm smears were prepared by placing 10 μl of each sperm suspension on cleaned microscopic slides and acrosome reacted spermatozoa were determined under a fluorescent microscope (Olympus Corporation, Japan) with excitation of 490 nm by counting 100 spermatozoa in 3 microscopic fields.

A spermatozoon is considered to be acrosome reacted if initial fusion phase or loss of acrosomal content was observed.

2.8 Determination of DNA damage in spermatozoa

Spermatozoa incubated with respective doses of tebufenozide and the control (BWW) for 1 h were used (n=6) for assessment of DNA damage. Sperm smears were prepared on cleaned microscopic slides for both treated and control samples and air dried for 5 minutes. The smears were fixed in Carnoy's solution for at least 3h. Slides were washed with distilled water, air-dried and stained with acridine orange for five minutes. Smears were evaluated with the aid of fluorescent microscope (Olympus Corporation, Japan) with excitation of 490 nm. Hundred sperm from each staining protocol were scored and graded. All sperm exhibiting yellow to red colour was scored as denatured DNA and sperm exhibiting green colour was scored as normal DNA (Tajeda et al. 1994).

2.9 Statistical analysis

Statistical analysis was performed with Two-way analysis of variance (ANOVA) using Minitab software package (Minitab Co., USA) for the main effect of pesticides. Where a significant treatment effect was found, differences among individual group means were tested by "Tukey 95%". The data are expressed as mean \pm SD. The p value was set to $P < 0.05$.

3 Results

3.1 Effects on sperm motility

According to figure 1, tebufenozide significantly ($P < 0.01$) decreased the motility at all dose levels respectively by 33.7%, 36.4% and 46.3% at 1.25, 2.5 and 5 $\mu\text{g/ml}$.

3.2 Sperm viability

Though sperm viability was decreased (figure 2) in spermatozoa treated with 2.5 and 5 $\mu\text{g/ml}$ doses of tebufenozide, a significant ($P < 0.05$) reduction of sperm viability (by 46.2 %) was recorded only in the highest dose.

3.3 Effects on DNA damage in spermatozoa

Percentage of DNA damaged spermatozoa count was increased at all tebufenozide treated doses. The percentages of damages were significantly ($p < 0.05$) increased by 164.8% with the lowest dose, by 326.9% and significantly increased ($p < 0.01$) at the mid dose and by 357.9% at the highest dose. Results are summarized in Table I.

3.4 Effects on sperm capacitation and sperm acrosome reaction

Human sperm incubated *in vitro* with tebufenozide elicited significant ($p < 0.05$) effects on sperm capacitation at all dose levels (by 20.6%: low dose; 45.5%: mid dose and 58.0%: high dose). Acrosome reaction was impaired only at the highest dose (36.6%). Results are summarized in Table 1.

4 Discussion

Results obtained from the present study indicated that tebufenozide could exert significant effect on human sperm *in vitro*. Similar results were observed with chlorinated hydrocarbon (Pflieger-Bruss & Schill 2010), methamidophos (Peiris et al., 1995) and Judo 40 (Kudavidanage and Peiris, 2016). In the present study, sperm motility was reduced in all dose levels of tebufenozide treated human sperms. Changes in mitochondrial membrane potential by different pesticides can result (Pant et al. 2014) and oxidative stress induced by organophosphorus pesticides (Lukaszewicz-Hussain 2010) could damage mitochondrial membrane to decrease the production of ATP thus affecting the motility (Wang et al. 2003). Disulfide bonds in the tail are vital for rigidity of spermatozoa tail that is essential for progressive motility and oxidation of disulfide bonds could result in loss of sperm motility (Buffon et al. 2012). Further, acridine orange test revealed that tebufenozide could alter integrity of the sperm head at both dose levels and hence it is possible that the same alteration could be observed in the disulfide bonds of the tail resulting impaired motility.

Reactive oxygen species could cause membrane damage resulting in continuous decrease in sperm motility and vitality after ejaculation (Calamera et al., 2001). Tebufenozide have been shown to cause oxidative damages (Carter et al., 2001) resulting in reduction of sperm motility, and viability. The Viability is the proportion of live spermatozoa determined by the evaluation of cellular and or membrane integrity

(Rao, 2006). Vitality of sperms associated with intact, functional and semipermeable plasma membranes (Mandani *et al.*, 2013). Disruption of DNA occurs mainly through the oxidation of disulfide bonds. It has been reported that high levels of ROS mediate the oxidation of disulfide bonds, which are commonly observed in the spermatozoa of infertile men (Aitken and Krausz 2001). Furthermore, it has been shown that pesticides are known to generate ROS thus producing DNA damage observed in the present study (Choi *et al.* 2015). Damage to disulfide bonds in the tail could also lead to reduction in capacitation, which was observed in the present study. Hyperactivation is an indication of sperm completion of capacitation and is essential for penetration through the cumulus mass and zona pellucida (Peiris and Moore, 2001a). Acrosome reaction is essential for successful penetration of the zona pellucida (Flesch and Gadella, 2000) and tebufenozide interfered with acrosome reaction. Tebufenozide could alter calcium ion concentrations, thus affecting spontaneously induced human spermatozoa acrosome reaction (Luconi *et al.*, 2001). Plasma membrane of sperm contains high levels of polyunsaturated fatty acids, which make spermatozoa susceptible to oxidative damage (Gravance *et al.* 2003). Hence alteration observed in the acrosome reacted spermatozoa probably due to membrane changes exerted by the insecticide (Aitken and Krausz, 2001).

5 Conclusion

From the results, it can be concluded that human sperm functions were decreased with exposure of tebufenozide. With longer exposure to tebufenozide, the risk of infertility may increase.

References

1. Aitken R.J., Krausz C. (2001). Oxidative stress, DNA damage and the Y chromosome. *Reproduction*, 122, 497-506.
2. Buffone M. G., Ijiri T. W., Cao W., Merdushev T., Aghajanian H. K. and Gerton G. L. (2012) Heads or tails? Structural events and molecular mechanisms that promote mammalian sperm acrosomal exocytosis and motility. *Molecular Reproduction Development*, 79, 4–18
3. Calamera J.C., Fernandez P.J. Buffone M.G., Acosta A.A. and Dancel G.F. (2001). Effect of long term *in vitro* incubation of human spermatozoa: functional parameters and catalase effect. *Andrologia*, 33:79–86.

4. Carter C.S, Ramsey M.M., Ingram R.L., Cashion A.B., Cefalu W.M., Wang Z.Q. and Sonntang W.E. (2001). Models of Growth Hormone and IGF-1 Deficiency Applications to Studies of Aging Processes and Life-Span Determination. *The Journal of Gereontology*, 53:8177-8188.
5. Choi H., Han C., Jin S., Kwon J.T., Jeong J., Kim J., Ham S., Jeon S., Joon Y. and Cho C. (2015). Reduced fertility and altered epididymal and sperm integrity in mice lacking ADAM7. *Biology of Reproduction*, 93: DOI\10.1095/biolreprod.115.130252.
6. Dharmadasa M., Nagalingam T. and Seneviratne P.H.M. (2008). Identification And Screening Of New Generation Insecticides Against Cardamom Thrips (*Sciothrips Cardamomi*) In Cardamom Cultivations In Sri Lanka. *Ceylon Journal of Science (Biological Science)*, 37: 137-142.
7. Flesch F.M. and Gadella, B.M. (2000). Dynamics of the mammalian sperm plasma membrane in the process of fertilization. *Biochemical and Biophysics Acta*, 1469: 197-235.
8. Gavance C.G., Garner D.L., Miller M.G. and Berger, T. (2003). Flow cytometric assessment of changes in rat sperm mitochondrial function after treatment with pentachlorophenol. *Toxicology In Vitro*, 17: 253-257.
9. Gupta D.S., Mills C.L. and Fraser L.R. (1993). Ca^{2+} related changes in the capacitation state of human spermatozoa assessed by a chlortetracycline fluorescence assay. *Journal of Reproduction and Fertility*, 99: 135-143.
10. Kudavidanage E.P. and Peiris D. C. (2016). Exposure of Judo 40 alters DNA integrity and sperm function of rat and human spermatozoa. *EPRA International Journal of Multidisciplinary Research*, 2: 12-19.
11. Luconi M., Bonaccorsi L., Forti G. and Baldi E. (2001). Effects of estrogenic compounds on human spermatozoa: evidence for interaction with a nongenomic receptor for estrogen on human sperm membrane. *Molecular and Cellular Endocrinology*, 178: 39-45.
12. Lemarie G., Terouanne, B., Mauvais P., Mitchel S. and Rahmani, R. (2004). Effects of organochlorine pesticides on human androgen receptor activation in vitro. *Toxicology and Applied Pharmacology*, 196, 235-246.

13. Lukaszewicz-Hussain A. (2010). Role of oxidative stress in organophosphate insecticide toxicity - Short review. *Pest Biochemistry and Physiology* 98, 145-150
14. Luconi M., Bonaccorsi L, Forti G. and Baldi E.(2001). Effects of estrogenic compounds on human spermatozoa: evidence for interaction with a nongenomic receptor for estrogen on human sperm membrane. *Molecular and Cellular Endocrinology* 178: 39–45. (doi:10.1016/S0303-7207(01)00416-6)
15. Mandani P., Desai K. and Highland H. (2013) Cytotoxic effects of Benzene metabolites on human sperm function: An in vitro study. *ISRN Toxicology*, 2013: Article ID: 397524.
16. Muangphra P., Tharapoom K., Euawong N., Namchote S. and Gooneratne, R. (2015). Chronic toxicity of commercial chlorpyrifos on earthworm *Pheretima peguana* *Environmental Toxicology*, 2015:doi 10.1002/tox.22150
17. Pant N., Shukla M., UpChaturvedi P.K., Saxena D.K., Gupta Y.K. and Adhyay A.D. (2014). Association between environmental exposure to p, p'-DDE and lindane and semen quality. *Environmental Science Pollution Research International*. 21:11009-11016.
18. Peiris L.D.C, Jayathunga Y.N.A. and Ratnasooriya W.D. (1995). Antireproductive effects in male rats exposed to methamidophos. *Ceylon Journal of Science*, 24: 53-9.
19. Peiris L.D.C. and Moore H.D.M. (2001a). Effects of acute and chronic doses of methoxy acetic acid on hamster sperm fertilizing ability. *Asian Journal of Andrology* 3: 209-216.
20. Peiris L.D.C. and Moore H.D.M. (2001b). Effects of acute and chronic doses of methoxy acetic acid on hamster sperm fertilizing ability. *Asian Journal of Andrology* 3: 185-191.
19. Perry M.J., Vennersb S.A., Barr D.B and Xu X. (2007) Environmental pyrethroid and organophosphorus insecticide exposures and sperm concentration. *Reproductive Toxicology*, 23:113–118.
20. Pflieger-Bruss S. and Schill W.B. (2010). Effects of chlorinated hydrocarbons on sperm function *in vitro*. *Andrologia*, 32: 311-315.

21. Rao K.A. (2006). The infertility Manual. (Eds. P.R. Brinsden & A.H. Sathananthan) pp. 123-145, Jaypee brothers' Medical publishers, India.
22. Ratnasooriya W.D., Jayatunga Y.N.A and Peiris L.D.C.(1996). Monocrotophos impairs the fertility of male rats. *Medical Science Research*,24: 403 406.
23. Recio-Vega R., Ocampo-Gomez G., Borja-Aburto V.H., Moran-Martinez J. and Cebrian-Garcia M.E. (2008). Organophosphorous pesticide exposure decreases sperm quality: association between sperm parameters and urinary pesticide levels. *Journal of Applied Toxicology*, 28:674–680.
24. Sankoh A.I., Whittle R., Semple K.T., Jones K.C. and Sweetman A.J. (2016). An assessment of the impacts of pesticide use on the environment and health of rice farmers in Sierra Leone. Environment International, e-publication ahead, doi.org/10.1016/j.envint.2016.05.034: 1-9.
25. Tejada R., Mitchell J.C., Normann A., Marik J. J. and Friedman S. (1994). A test for the practical evaluation of male fertility by acridine orange fluorescence. *Fertility and Sterility*, 42: 87- 89.
26. Wang X., Sharma R.K., Gupta A., George V., Thomas A.J Jr., Falcone T. and Agarwal A. (2003). Alterations in mitochondria membrane potential and oxidative stress in infertile men: a prospective observational study. *Fertility and Sterility*, 80: 844–850.




Figure 1: The effects of different doses of tebufenozide (1.25 µg/ml; 2.5 µg/ml and 5 µg/ml) or BWW (control) on sperm motility. Results are presented as mean \pm SE Mean (n=6). **P \leq 0.01 significantly different from control.




Figure 2: Effects of different concentrations of tebufenozide (1.25 µg/ml, 2.5 µg/ml and 5 µg/ml) and BWW (control). Results are presented as mean \pm SE Mean (n=9). *P \leq 0.05, significantly different from control.

Table I. Effects of tebufenozide on human sperm DNA damage, capacitation and acrosome reaction

Parameters	Control (BWW)	Treatment (µg/ml) (Tebufenozide)		
		1.25	2.5	5
DNA damaged spermatozoa (%)	14.5 \pm 0.03	38.4 \pm 0.04*	61.9 \pm 0.25**	80.9 \pm 1.15**
Capacitated spermatozoa (%)	68.6 \pm 0.94	54.5 \pm 0.65*	37.4 \pm 0.89**	28.8 \pm 1.0**
Acrosome reacted spermatozoa (%)	80.9 \pm 0.02	70.8 \pm 0.56	71.6 \pm 0.47	51.3 \pm 0.91**

The data are given as mean \pm SEM (n=6). Values are statistically significant at *p<0.05, ** p < 0.01. Control group was treated with BWW while treated groups included 1.25, 2.5 & 5 µg/ml of tebufenozide.